

学会名	開催都市名	題名
Spine World Summit	Hongkong, January, 2018.	Clinical and radiological outcome of microendoscopic laminotomy for degenerative lumbar spondylolisthesis –In comparison with the cases without spondylolisthesis Balloon kyphoplasty for fresh osteoporotic vertebral fractures with poor prognostic factors – A multicenter prospective intervention study –
American Academy of Orthopaedic Surgeon	New Orleans, March, 2018.	Peri-prosthetic Bone Mineral Density (BMD) after Simultaneously Bilateral Total Can Porous Tantalum Tibial Component Prevent Periprosthetic Loss of Bone Mineral Density After Total Knee Arthroplasty More Than Five Years? Does RA patient have lumbar lesion or low back pain more frequently than healthy population? –Cross sectional analysis in cohort study with age and sex matched healthy volunteer Differences in clinical and radiological outcomes depending on timing of balloon kyphoplasty for painful osteoporotic vertebral fracture Impact of Hemodialysis on surgical outcomes and mortality rate after lumbar spine surgery – a matched cohort study –
Orthopedic Research Society	New Orleans, March, 2018.	The degeneration of adjacent intervertebral discs negatively influences union rate of osteoporotic vertebral fracture: A multicenter cohort study Peri-prosthetic Bone Mineral Density (BMD) after Simultaneously Bilateral Total Knee Arthroplasty (TKA) Under Oral Alendronate Therapy 2D Measurement Misidentifies the Alignment Outliers in Total Knee Arthroplasty Comparison of the Periprosthetic Bone Mineral Density Change between Medial Pivot PS Prosthesis and Conventional PS Prosthesis at Three Years A Mobile-Bearing Total Knee Arthroplasty Prevents Periprosthetic Loss of Bone Mineral Density Around the Femoral Component for 10 Years Can Porous Tantalum Tibial Component Prevent Periprosthetic Loss of Bone Mineral Density After Total Knee Arthroplasty? –A Mean of 10 Years Follow Up– Joint Line Elevation Did Not Induce the Mid-Flexion Joint Gap Laxity in Total Knee Arthroplasty The Risk of Patella-post Impingement Differed between the Tibial Post Designs of Posterior Stabilized Prostheses A New Patient Specific Instrumentation Improved the Accuracy of Tibial Rotation for Total Knee Arthroplasty
18th ESSKA	Glasgow, May, 2018.	Donor site evaluation after osteochondral autograft transplantation for capitellar Prevalence and Risk Factors for Meniscal Cyst Due to Meniscal Repair Changes of the residual meniscus after reshaping surgery for discoid lateral Tegner Activity Scale as Predictor of Re-injury after Double-bundle Anterior Cruciate Ligament Reconstruction in Young Athletes

		<p>Concentrations of serum cartilage oligomeric matrix protein before and after anterior cruciate ligament reconstruction.</p> <p>Histological analysis of femoral attachment of Anterior Cruciate Ligament</p> <p>The relationship between tibial tunnel placement and graft intensity on MRI in anatomical double-bundle ACL reconstruction.</p>
45th Annual Meeting of International Society for the Study of Lumbar Spine	Canada,May,2018.	<p>EFFICACY OF BALLOON KYPHOPLASTY FOR FRESH OSTEOPOROTIC VERTEBRAL FRACTURES WITH POOR PROGNOSTIC FACTORS – A PROPENSITY-SCORE-MATCHED ANALYSIS USING DATA FROM THE MULTICENTER PROSPECTIVE STUDY –</p> <p>Differences in clinical and radiological outcomes depending on timing of balloon kyphoplasty for painful osteoporotic vertebral fracture</p> <p>Factors related to spinal sagittal imbalance among an elderly population in a suburban community: The Shiraniwa study</p>
The 19th European Federation of National Associations of Orthopaedics and Traumatology	Barcelona,May,2018.	<p>Ultrasound Imaging Diagnosis Of Soft Tissue Tumors Based On WHO Classification.</p> <p>Prognostic Factors For Elderly Patients With Primary Malignant Bone And Soft Tissue Tumors</p> <p>Diagnostic Value Of Tumor-Fascia Relationship In Superficial Soft Tissue Masses On Magnetic Resonance Imaging</p> <p>Accuracy Of Core Needle Biopsy For Soft Tissue Tumors</p>
5th APKASS	Sydney,May,2018.	<p>Prognostic Factors of Clinical Failure of The Osteochondral Autograft Transplantation for Lateral Wide Spread Type of Unstable Osteochondral Dissecans of Humeral Capitellum</p>
EULAR2018	Amsterdam,June,2018	<p>Matrix metalloproteinase-3 is a good predictor for joint destruction only in male patients with rheumatoid arthritis.</p> <p>Comparison of clinical features in patients with psoriatic arthritis and patients with spondyloarthritis with inflammatory bowel disease</p> <p>Ultrasonographic assessment of nail in psoriatic disease found a link between nail disorder and soft tissue swelling around the nail</p> <p>Severity of skin symptoms is not associated with musculoskeletal manifestations in patients with psoriasis</p> <p>Comparison of bone mineral density between rheumatoid arthritis patients and healthy individuals over seven years from the tomorrow study</p> <p>The comparison of the ultrasonographic synovial findings between intravenous administration and subcutaneous injection of tocilizumab</p> <p>Effect of aging on bone mass and skeletal muscle mass in elderly patients with rheumatoid arthritis: tomorrow study</p>

		<p>Screening research of ultrasonographic peripheral arthritis and enthesitis in patients with inflammatory bowel disease</p> <p>Increased homocysteine level for 7 years in patients with rheumatoid arthritis: tomorrow study</p> <p>Glucocorticoid use is an independent risk factor for sarcopenia in patients with rheumatoid arthritis – from the chikara study –</p> <p>The effect of concomitant type of vitamin d, biological dmards and disease activity for therapeutic effect of denosumab in osteoporosis patients with rheumatoid</p>
Federation of European Societes for Surgery of the Hand	Copenhagen,June,2018.	Low second to fourth digit ratio in Dupuytren disease.
Federation of European Societes for Surgery of the Hand	Copenhagen,June,2018.	Radial Artery Perforator Adiposal Flap for Coverage of the Neurolysed Median Nerve
The 18th Annual Meeting Pacific and Asian Society of Minimally Invasive Spine Surgery	Kaohsiung,July,2018.	<p>The Incidence of Nerve Root Injury by Highspeed Drill Can be Reduced by Chilled Saline Irrigation in a Rabbit Model</p> <p>Anterior Corpectomy and Reconstruction using X-CORE 2 VBR SYSTEM® for Osteoporotic Vertebral Fractures</p> <p>Clinical impact of Sarcopenia for the Outcomes of Minimally Invasive Lumbar Decompression Surgery</p> <p>Clinical result of CT-guided percutaneous vertebroplasty for metastatic spinal</p> <p>Does teriparatide impact the clinical and radiographic outcome of balloon kyphoplasty for fresh osteoporotic vertebral fracture?</p> <p>Differences in clinical and radiological outcomes depending on timing of balloon kyphoplasty for painful osteoporotic vertebral fracture</p>
Spine Across the Sea 2018	Hawaii,July,2018.	<p>Balloon kyphoplasty for fresh osteoporotic vertebral fractures with poor prognostic factors</p> <p>– Multicenter prospective intervention study –</p> <p>The degeneration of adjacent intervertebral discs negatively influences union rate of osteoporotic vertebral fracture: A multicenter cohort study</p> <p>Preoperative severity of facet joint degeneration does not impact on 2-year clinical outcomes following laminoplasty</p> <p>Thoracic Spine Translation and Angular Motion: an Analysis using Thoracic Spine Kinematic MRI (kMRI)</p> <p>Factors related to spinal sagittal imbalance among an elderly population in a Japanese suburban community: The Shiraniwa study</p>
The 5th Tissue Engineering and Regenerative Medicine International Society World Congress	Kyoto,September,2018.	Dual release of bFGF and SDF-1 accelerates peripheral nerve regeneration.

73rd American Society for Surgery of the Hand	Boston,September,2018.	Arthroscopic Hemitrapeziectomy with Suspensionplasty using a Half Slip of Extensor Carpi Radialis Longus Tendon for Treatment of Thumb Carpometacarpal Arthritis
EURO SPINE 2018	Barcelona,September,2018.	Balloon kyphoplasty versus conservative treatment for acute osteoporotic vertebral fractures with poorprognostic factors – An Propensity–Score–Matched Analysis Using Data from the Two Prospective Multicenter studies –
		Related factors of Neck Disability Index (NDI) among elderly people in suburban area: Shiraniwa cohort study
		Does teriparatide impact the clinical and radiographic outcome of balloon kyphoplasty for fresh osteoporotic vertebral fracture?
39th SICOT Orthopaedic World Congress	Montréal,October,2018.	Diagnostic imaging method using magnetic resonance imaging of superficial soft tissue masses
ACR2018	Chicago,Octorber,2018.	Adipose Derived Stem Cell Suppressed Synovial Inflammation and Repaired Cartilage Destruction in Rheumatoid Arthritis Model Mice
		Ultrasonographic Research of the Relationship between Nail Disorders and Peripheral Arthritis or Enthesitis in Patients with Psoriasis and Psoriatic Arthritis
Cervical Spine Research Society	Scottsdale,December, 2018.	The Predictor Of Patients Who Failed To Achieve The Minimum Clinical Important Differences Following Laminoplasty For Cervical Spondylosis Myelopathy